PRODUCT INFORMATION



17(R)-HDHA

Item No. 10005099

CAS Registry No.:	155976-53-9
Formal Name:	17R-hydroxy-
	4Z,7Z,10Z,13Z,15E,19Z-
	docosahexaenoic acid
Synonyms:	17(R)-hydroxy Docosahexaenoic Acid, COOH
	17(R)-HDoHE
MF:	C ₂₂ H ₃₂ O ₃
FW:	344.5 OH
Purity:	≥98%
Stability:	≥2 years at -20°C
Supplied as:	A solution in ethanol
UV/Vis.:	λ _{max} : 236 nm ε: 10,000

Laboratory Procedures

For long term storage, we suggest that 17(R)-HDHA be stored as supplied at -20°C. It should be stable for at least two years.

17(R)-HDHA is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. 17(R)-HDHA is miscible in these solvents.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 17(R)-HDHA is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of 17(R)-HDHA in PBS (pH 7.2) is approximately 0.8 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

Resolvins are a group of polyhydroxylated metabolites of docosahexaenoic acid (DHA) found in the inflammatory exudates of aspirin-treated experimental animals.¹ 17(R)-HDHA is the primary oxygenation product of DHA when exposed to aspirin-inhibited cyclooxygenase-2. 17(R)-HDHA serves as a precursor to resolvins and has intrinsic biological activity, such as the inhibition of TNF α -induced IL-1 β expression in human glioma cells and inhibition of TNF α -induced leukocyte trafficking to the murine air pouch.¹

Reference

1. Serhan, C.N., Hong, S., Gronert, K., et al. Resolvins: A family of bioactive products of .omega.-3 fatty acid transformation circuits by aspirin treatment that counter proinflammation signals. J. Exp. Med. 196(8), 1025-1037 (2002).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/25/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM