PRODUCT INFORMATION



SB-743921 (hydrochloride)

Item No. 18384

CAS Registry No.:	940929-33-9	`
Formal Name:	N-(3-aminopropyl)-N-[(1R)-1-[7-	
	chloro-4-oxo-3-(phenylmethyl)-4H-1-	0
	benzopyran-2-yl]-2-methylpropyl]-4-	
	methyl-benzamide, monohydrochloride	
MF:	$C_{31}H_{33}CIN_2O_3 \bullet HCI$	
FW:	553.5	<pre></pre>
Purity:	≥98%	0
UV/Vis.:	λ _{max} : 203, 309 nm	
Supplied as:	A crystalline solid	
Storage:	-20°C	
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

SB-743921 (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the SB-743921 (hydrochloride) in the solvent of choice. SB-743921 (hydrochloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of SB-743921 (hydrochloride) in ethanol is approximately 20 mg/ml and approximately 50 mg/ml in DMSO and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of SB-743921 (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of SB-743921 (hydrochloride) in PBS, pH 7.2, is approximately 0.2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

SB-743921 is a potent inhibitor of the kinesin spindle protein Eg5 ($K_i = 0.5$ nM for microtubule-stimulated ATPase activity).¹ It inhibits the growth of HCT116 colon, LNCaP and PC3 prostate, K562 leukemia, BxPC-3 pancreas, and NCI-H1299 lung cancer cell lines in vitro (GI₅₀s = 25, 22, 48, 50, 80, and 82 nM, respectively). In vivo, SB-743921 induces complete tumor regression in a COLO 205 mouse xenograft model and partial regression in MCF-7, SK-MES, H69, and OVCAR-3 mouse xenograft models.² SB-743921 (2.5-10 mg/kg) also reduces tumor volume in a dose-dependent manner in a diffuse large B cell lymphoma (DLBCL) mouse xenograft model.³

References

- 1. Good, J.A.D., Wang, F., Rath, O., et al. J. Med. Chem. 56(5), 1878-1893 (2015).
- 2. Jackson, J.R., Gilmartin, A., Dhanak, D., et al. Clin. Cancer Res. 12(19 Suppl) (2006).
- 3. Bongero, D., Paoluzzi, L., Marchi, E., et al. Leuk. Lymphoma 56(10), 2945-2952 (2015).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/09/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM