

PRODUCT INFORMATION



Mocetinostat

Item No. 18287

CAS Registry No.: 726169-73-9

Formal Name: N-(2-aminophenyl)-4-[[[4-(3-pyridinyl)-2-pyrimidinyl]amino]methyl]-benzamide

Synonym: MGCD0103

MF: C₂₃H₂₀N₆O


FW: 396.4

Purity: ≥98%

Stability: ≥2 years at -20°C

Supplied as: A crystalline solid

UV/Vis.: λ_{max}: 237 nm



Laboratory Procedures

For long term storage, we suggest that mocetinostat be stored as supplied at -20°C. It should be stable for at least two years.

Mocetinostat is supplied as a crystalline solid. A stock solution may be made by dissolving the mocetinostat in the solvent of choice. Mocetinostat is soluble in organic solvents such as DMSO and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of mocetinostat in these solvents is approximately 20 and 25 mg/ml, respectively.

Mocetinostat is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, mocetinostat should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Mocetinostat has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Mocetinostat is an orally available inhibitor of histone deacetylases (HDACs) that selectively targets HDAC1 and 2 (IC₅₀s = 0.15 and 0.29 μM, respectively), less potently inhibits HDAC3 and 11 (IC₅₀s = 1.66 and 0.59 μM, respectively) and has negligible actions against HDAC4-8.^{1,2} It induces hyperacetylation of histones, causes cell cycle blockade, and induces apoptosis in cancer cells *in vitro*.^{1,3} Mocetinostat also has antitumor activity *in vivo*, blocking the growth of human tumor xenografts in mice.¹ It also significantly reduces cardiac fibrosis following ischemic heart failure in mice.⁴

References

1. Fournel, M., Bonfils, C., Hou, Y., *et al.* MGCD0103, a novel isotype-selective histone deacetylase inhibitor, has broad spectrum antitumor activity *in vitro* and *in vivo*. *Mol. Cancer Ther.* **7**(4), 759-768 (2008).
2. Marson, C.M., Matthews, C.J., Atkinson, S.J., *et al.* Potent and selective inhibitors of histone deacetylase-3 containing chiral oxazoline capping groups and a N-(2-aminophenyl)-benzamide binding unit. *J. Med. Chem.* **58**, 6803-6818 (2015).
3. Bonfils, C., Kalita, A., Dubay, M., *et al.* Evaluation of the pharmacodynamic effects of MGCD0103 from preclinical models to human using a novel HDAC enzyme assay. *Clin. Cancer Res.* **14**(11), 3441-3449 (2008).
4. Nural-Guvener, H., Zakharova, L., Feehery, L., *et al.* Anti-fibrotic effects of class I HDAC inhibitor, mocetinostat is associated with IL-6/Stat3 signaling in ischemic heart failure. *Int. J. Mol. Sci.* **16**, 11482-11499 (2015).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/10/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM