

TRIP12 Polyclonal Antibody

| Catalog No. | A9958 | Category | Polyclonal Antibodies | |
|--|---|---------------|----------------------------------|--|
| Applications | WB, IHC, IF | Observed MW | 260kDa | |
| Cross-reactivity | Human, Mouse, Rat | Calculated MW | 192kDa/220kDa/223kDa/225kDa | |
| Immunogen Information | | | Recommended Dilutions | |
| Immunogen | Recombinant fusion protein containing corresponding to amino acids 1-240 of (NP_001271143.1). | | WB 1:500 - 1:1000 | |
| Gene ID | 9320 | | IHC 1:50 - 1:100 | |
| Swiss prot | Q14669 | | | |
| Synonyms | TRIP12; TRIP-12; ULF; E3 ubiquitin-prote ase TRIP12 | ein lig | IF 1:50 - 1:100 | |
| Product information | | | 300kDa - | |
| Source | Rabbit | | 250kDa — — TRIP12 | |
| lsotype | IgG | | 180kDa — | |
| Purification method | Affinity purification | | | |
| Storage | Store at -20°C. Avoid freeze / thaw cycl Buffer: PBS with 0.02% sodium azide, 5 | | 130kDa — | |
| Background | | | 100kDa — | |
| The protein encoded by this gene is an E3 ubiquitin-protein ligase involved in the | | | Western blot - TRIP12 Polyclonal | |

The protein encoded by this gene is an E3 ubiquitin-protein ligase involved in the degradation of the p19ARF/ARF isoform of CDKN2A, a tumor suppressor. The encoded protein also plays a role in the DNA damage response by regulating the stability of USP7, which regulates tumor suppressor p53.

