

Substrates for Biochemical Assays

For Alkaline Phosphatase

4-Nitrophenyl Phosphate (Ready-to-use solution) [for ELISA]
(= pNPP (Ready-to-use solution))

100mL [N1109]

Application

1. Add 100 μ L of pNPP solution (Product No. N1109) to each well.
2. Incubate the plate at room temperature for 30 minutes.
3. To terminate the reaction, add 100 μ L of 1N NaOH solution (Product No. S0542) to each well.
4. Within 1 hour from start the reaction, measure the absorbance of each well at 405 nm.




Figure. An example of use by the above method

Soluble Substrates

4-Nitrophenyl Phosphate Disodium Salt Hexahydrate (= pNPP)	1g / 5g [D4005]
4-Nitrophenyl Phosphate Di(tris) Salt Hydrate	5g / 25g [N0422]
1-Naphthylphosphoric Acid Monosodium Salt Monohydrate	1g / 5g / 25g [N0452]
1-Naphthylphosphoric Acid Disodium Salt Hydrate	1g / 5g [P0263]

Precipitate Substrates

For such as immunohistochemical staining or immunoblotting, substrates arising precipitate dyes with alkaline phosphatase.

Fast Blue RR Salt	5g / 25g [B0785]
X-Phosphate <i>p</i>-Toluidine Salt	100mg / 1g [B1239]
Blue Tetrazolium	1g / 5g [B3581]
Naphthol AS-TR Phosphate	200mg [C2250]
Nitro Blue Tetrazolium (= NBT)	100mg / 1g [D0844]
Fast Black K Salt	5g [F0734]
Iodonitrotetrazolium Chloride (= INT)	100mg / 1g [I0781]
Tetranitro Blue Tetrazolium (= TNBT)	100mg / 1g [T0250]

For Peroxidase

TMB [for ELISA] (Ready-to-use solution)

(= 3,3',5,5'-Tetramethylbenzidine (Ready-to-use solution))

100mL [T3854]

Application

1. Add 100 μ L of TMB solution (Product No. T3854) to each well.
2. Incubate the plate at room temperature for 30 minutes.
3. The reaction, add 100 μ L of 1N HCl solution (Product No. H1202) to each well.
4. Measure the absorbance of each well at 450 nm.

When this product T3854 reacts with horseradish peroxidase (HRP), a blue colored soluble reaction product appears thus it can be used for ELISA.

This product cannot be used for Western blotting which needs a precipitate.




Figure.
An example of use by the above method

TMB [for Western blotting] (Ready-to-use solution)

(= 3,3',5,5'-Tetramethylbenzidine (Ready-to-use solution))

100mL [T3855]

Application

1. Incubate a blotting membrane with an HRP-conjugated antibody and then wash the membrane.
2. Incubate the washed membrane with TMB solution (Product No. T3855) until color development.
3. Add deionized water to stop color development.

When this product T3855 reacts with HRP, a blue-purple precipitate appears thus it can be used for Western blotting.

This product cannot be used for ELISA which needs a soluble reaction product.



M 1

Figure.
An example of Western blotting by the above method

M : molecular weight marker
1 : Target protein A

4-CN (Ready-to-use solution) [for Western blotting]
(= 4-Chloro-1-naphthol (Ready-to-use solution))

100mL [C3384]

Application

1. Incubate a blotting membrane with an HRP-conjugated antibody and then wash the membrane.
2. Incubate the washed membrane with 4-CN solution (Product No. C3384) until color development.
3. Add deionized water to stop color development.




Figure.
An example of Western blotting by the above method

M : molecular weight marker
1 : Target protein B (Middle concentration)
2 : Target protein B (Low concentration)

AzBTS (Ready-to-use solution) [for ELISA]

(= 2,2'-Azinobis(3-ethylbenzothiazoline-6-sulfonic Acid Ammonium Salt) (Ready-to-use solution))

100mL [A3176]

Application

1. Add 100 μ L of AzBTS solution (Product No. A3176) to each well.
2. Incubate the plate at room temperature for 30 minutes.
3. Within 1 hour from start the reaction, measure the absorbance of each well at 405 nm.




Figure.
An example of use by the above method

DAB staining kit

1kit [D5909]

Advantage

DAB substrate solution is used for detecting HRP-conjugated antibodies by Western blot analysis. Product No. **D5909** includes three kinds of reagents which are necessary for preparing the DAB substrate solution. The substrate solution is easily prepared by mixing each component. This kit includes reagents for 100 tests (about 500 mL) of the DAB substrate solution.



Application

1. Prepare a blotting membrane covered with HRP-conjugated antibody.
2. Prepare 5mL of deionized water, add these solutions in the kit and prepare the DAB substrate solution.
 - Add 5 drops of DAB solution (about 140 μ L)
 - Add 2 drops of Hydrogen Peroxide solution (about 80 μ L)
 - Add 2 drops of 1.5M Tris-HCl buffer (about 80 μ L)
3. Soak the membrane in the DAB substrate solution and incubate until the membrane is colored.
4. Add deionized water and stop the coloring reaction.

Figure.
An example of Western blotting by the above method

M : molecular weight marker
1 : Target protein C



Related Products

Sodium Hydroxide (1mol/L in Water)	500mL [S0542]
Hydrochloric Acid (1mol/L)	500mL [H1202]
Peroxidase from Horseradish	100mg / 1g [P0073]
Horseradish Peroxidase Maleimide Conjugate (0.5mg \times 3)	1set [H1621]
Horseradish Peroxidase NHS Ester Conjugate (0.2mg \times 3)	1set [H1746]
Anti-6xHis Monoclonal Antibody (6A12) HRP Conjugate	0.05mg/1vial [A3075]
Anti-Protein A Chicken Polyclonal Antibody HRP Conjugate	0.05mg/1vial [A3187]
Anti-αGal Chicken Polyclonal Antibody HRP Conjugate	0.05mg/1vial [A3195]
Anti-NeuGc Polyclonal Antibody HRP Conjugate	0.05mg/1vial [A3397]
Goat Anti-Mouse IgG HRP Conjugate	0.1mg/1vial [G0407]
Goat Anti-Mouse IgM HRP Conjugate	0.1mg/1vial [G0417]
Goat Anti-Rabbit IgG HRP Conjugate	0.1mg/1vial [G0418]
Sheep Anti-Chicken IgY HRP Conjugate	0.1mg/1vial [S0999]
Protein A HRP Conjugate	0.2mg/1vial [P2466]
Streptavidin HRP Conjugate	0.1mg/1vial [S0972]

Soluble Substrates (for ELISA etc.)

For such as ELISA, substrates generating soluble dyes with peroxidase.

AzBTS (= 2,2'-Azinobis(3-ethylbenzothiazoline-6-sulfonic Acid Ammonium Salt))	1g	[A2166]
OPD·2HCl (= 1,2-Phenylenediamine Dihydrochloride)	1g / 5g	[P1144]
OPD (= 1,2-Phenylenediamine)	1g / 5g	[P1805]
TMB (= 3,3',5,5'-Tetramethylbenzidine)	1g / 5g	[T2573]

Soluble Substrates (for determining H₂O₂)

Substrates generating soluble dyes for determining hydrogen peroxidase (H₂O₂) by various enzyme reactions.

4-AA·2HCl (= 4-Aminoantipyrine Hydrochloride)	5g / 25g	[A0257]
4-AA (= 4-Aminoantipyrine)	1g / 5g	[A2254]
5-ASA (= 5-Aminosalicylic Acid) * ¹	5g / 25g	[A2291]
DCHBS (= 3,5-Dichloro-2-hydroxybenzenesulfonic Acid Sodium Salt) * ¹	25g	[D1928]
2,4-DCP (= 2,4-Dichlorophenol) * ¹	1g / 5g	[D3865]
DMA (= <i>N,N</i> -Dimethylaniline) * ¹	1g / 5g	[D3866]
DMT (= <i>N,N</i> -Diethyl- <i>m</i> -toluidine) * ¹	1g / 5g	[D3868]
TOOS Hydrate (= Sodium 3-[Ethyl(<i>m</i> -tolyl)amino]-2-hydroxy-1-propanesulfonate Hydrate) * ¹	1g / 5g	[S0805]
ALPS (= Sodium 3-(<i>N</i> -Ethylanilino)propanesulfonate) * ¹	200mg / 1g	[S0817]
ADOS (= Sodium 3-(<i>N</i> -Ethyl-3-methoxyanilino)-2-hydroxy-1-propanesulfonate) * ¹	200mg / 1g	[S0826]
HDAOS (= <i>N</i> -(2-Hydroxy-3-sulfopropyl)-3,5-dimethoxyaniline Sodium Salte) * ¹	200mg / 1g	[S0827]
MBTH·HCl (= 3-Methyl-2-benzothiazolinonehydrazone Hydrochloride)	1g / 5g	[M2155]

*¹: Used together with [A2254](#) (or [A0257](#))

Precipitate Substrates

For such as immunohistochemical staining or immunoblotting, substrates arising precipitate products with peroxidase.

AEC (= 3-Amino-9-ethylcarbazole)	1g / 5g	[A2167]
4-CN (= 4-Chloro-1-naphthol)	1g / 5g	[C2291]
DAB (= 3,3'-Diaminobenzidine)	1g / 5g	[D3756]
DAB·4HCl (= 3,3'-Diaminobenzidine Tetrahydrochloride Hydrate)	1g / 5g	[D3757]
o-Dianisidine * ²	1g / 5g	[D3864]
o-Dianisidine Dihydrochloride * ²	1g / 5g	[D3893]
DMPD·2HCl (= <i>N,N</i> -Dimethyl-1,4-phenylenediamine Dihydrochloride) * ³	1g / 5g	[D3931]
1-Naphthol * ³	1g / 5g	[N0864]

*²: By combining [N0864](#) and [D3931](#) *³: Used together with [C2291](#)

For β -Galactosidase

Generating Insoluble Dyes

X-Gal (= 5-Bromo-4-chloro-3-indolyl β -D-Galactopyranoside) ■ Blue	200mg / 1g [B3201]
Magenta-Gal (= 5-Bromo-6-chloro-3-indolyl β -D-Galactopyranoside) ■ Purplish red	20mg / 100mg [B3469]
Bluo-Gal (= 5-Bromo-3-indolyl β -D-Galactopyranoside) ■ Deep blue	20mg / 100mg [B3470]
Salmon-Gal (= 6-Chloro-3-indolyl β -D-Galactopyranoside) ■ Light purplish red	20mg / 100mg [C2371]

Generating Soluble Dyes

ONPG (= 2-Nitrophenyl β -D-Galactopyranoside) ■ Yellow	1g / 5g / 25g [N0418]
PNPG (= 4-Nitrophenyl β -D-Galactopyranoside) ■ Yellow	1g / 5g [N0616]

For β -Glucuronidase

X-Gluc CHA Salt (= 5-Bromo-4-chloro-3-indolyl β -D-Glucuronide Cyclohexylammonium Salt) ■ Blue	10mg / 100mg [B3620]
X-Gluc Sodium Salt (= 5-Bromo-4-chloro-3-indolyl β -D-Glucuronide Sodium Salt) ■ Blue	10mg / 100mg [B3621]

For Luciferase

D-(-)-Luciferin	10mg / 50mg [A5030]
------------------------	---------------------

Every reagent in the category "Substrate" is manufactured for biochemical research.

For further information please refer to our website at www.TCIchemicals.com. ▶▶▶

Substrate



Ordering and Customer Service

TCI AMERICA

Tel : 800-423-8616 / 503-283-1681
Fax : 888-520-1075 / 503-283-1987
E-mail : Sales-US@TCIchemicals.com

TCI EUROPE N.V.

Tel : +32 (0)3 735 07 00
Fax : +32 (0)3 735 07 01
E-mail : Sales-EU@TCIchemicals.com

TCI Deutschland GmbH

Tel : +49 (0)6196 64053-00
Fax : +49 (0)6196 64053-01
E-mail : Sales-DE@TCIchemicals.com

Tokyo Chemical Industry UK Ltd.

Tel : +44 (0)1865 784560
Fax : +44 (0)1865 784561
E-mail : Sales-UK@TCIchemicals.com

TCI Chemicals (India) Pvt. Ltd.

Tel : 1800 425 7889 / 044-2262 0909
Fax : 044-2262 8902
E-mail : Sales-IN@TCIchemicals.com

梯希爱(上海)化成工业发展有限公司

Tel : 800-988-0390 / 021-67121386
Fax : 021-6712-1385
E-mail : Sales-CN@TCIchemicals.com

TOKYO CHEMICAL INDUSTRY CO., LTD.

Tel : +81 (0)3-5640-8878
Fax : +81 (0)3-5640-8902
E-mail : globalbusiness@TCIchemicals.com

Availability, price or specification of the listed products are subject to change without prior notice. Reproduction forbidden without the prior written consent of Tokyo Chemical Industry Co., Ltd.