

PRODUCT INFORMATION



Ro 01-6128

Item No. 11991

CAS Registry No.: 302841-86-7

Formal Name: N-(2,2-diphenylacetyl)-carbamic acid, ethyl ester

MF: C₁₇H₁₇NO₃


FW: 283.3

Purity: ≥98%

Supplied as: A crystalline solid

Storage: -20°C

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

Ro 01-6128 is supplied as a crystalline solid. A stock solution may be made by dissolving the Ro 01-6128 in the solvent of choice. Ro 01-6128 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of Ro 01-6128 in ethanol and DMSO is approximately 20 mg/ml and approximately 11 mg/ml in DMF.

Ro 01-6128 is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, Ro 01-6128 should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Ro 01-6128 has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Glutamate, the major excitatory neurotransmitter in the CNS, activates eight known subtypes of metabotropic glutamate receptors (mGluRs). Highly selective modulators designed to act at allosteric sites on certain mGluR subtypes are being developed to preferentially regulate subtype-specific, glutamate-induced receptor activation.¹ Ro 01-6128 is a positive allosteric modulator of mGluR1 that can potentiate glutamate-induced calcium release at rat mGluR1a with an EC₅₀ value of 104 nM.^{2,3} Ro 01-6128 activates ERK1/2 phosphorylation in the absence of exogenously added glutamate with an EC₅₀ value of 248 nM and potentiates glutamate-induced cAMP production with an EC₅₀ value of 21.5 μM.¹

References

1. Sheffler, D.J. and Conn, P.J. Allosteric potentiators of metabotropic glutamate receptor subtype 1a differentially modulate independent signaling pathways in baby hamster kidney cells. *Neuropharmacology* **55**(4), 419-427 (2008).
2. Knoflach, F., Mutel, V., Jolidon, S., et al. Positive allosteric modulators of metabotropic glutamate 1 receptor: Characterization, mechanism of action, and binding site. *Proc. Natl. Acad. Sci. USA* **98**(23), 13402-13407 (2001).
3. Hemstapat, K., de Paulis, T., Chen, Y., et al. A novel class of positive allosteric modulators of metabotropic glutamate receptor subtype 1 interact with a site distinct from that of negative allosteric modulators. *Mol. Pharmacol.* **70**(2), 616-626 (2006).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/10/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM