

PRODUCT INFORMATION



Arctiin

Item No. 15375

CAS Registry No.: 20362-31-6

Formal Name: (3R,4R)-4-[(3,4-dimethoxyphenyl)methyl]-3-[[4-(β-D-glucopyranosyloxy)-3-methoxyphenyl]methyl]dihydro-2(3H)-furanone

Synonyms: Arctigenin-4-Glucoside, NSC 315527

MF: C₂₇H₃₄O₁₁

FW: 534.6


Purity: ≥98%

UV/Vis.: λ_{max}: 229, 280 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

Arctiin is supplied as a crystalline solid. A stock solution may be made by dissolving the arctiin in the solvent of choice. Arctiin is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of arctiin in ethanol is approximately 2 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Arctiin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, arctiin should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Arctiin has a solubility of approximately 0.5 mg/ml in a 1:1 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Arctiin is the major active lignin in fruits of the burdock plant *A. lappa*. It demonstrates potent antiviral activity against influenza A virus and anti-inflammatory effects by decreasing the production of nitric oxide and pro-inflammatory cytokines.^{1,2} Arctiin is metabolized by human intestinal bacteria into various bioactive metabolites including arctigenin (Item No. 14913) and enterolactone (Item No. 10112), which respectively exhibit growth inhibitory and growth promoting activity in MCF-7 breast cancer cells at 10 μM.³

References

1. Hayashi, K., Narutaki, K., Nagaoka, Y., *et al.* Therapeutic effect of arctiin and arctigenin in immunocompetent and immunocompromised mice infected with influenza A virus. *Biol. Pharm. Bull.* **33(7)**, 1199-1205 (2010).
2. Lee, S., Shin, S., Kim, H., *et al.* Anti-inflammatory function of arctiin by inhibiting COX-2 expression via NF-κB pathways. *J. Inflamm. (Lond)* **8(1)**, 16 (2011).
3. Xie, L.-H., Ahn, E.-M., Akao, T., *et al.* Transformation of arctiin to estrogenic and antiestrogenic substances by human intestinal bacteria. *Chem. Pharm. Bull. (Tokyo)* **51(4)**, 378-384 (2003).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/26/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM