# **CelExplorer** Labs

## **Product Information**

# FocusClear<sup>TM</sup> Kit

A water-soluble clearing agent making biological tissue transparent

FocusClear <sup>TM</sup> MountClear <sup>TM</sup>

Immersion Solution-M

#### **1. INTRODUCTION**

Why using *FocusClear*<sup>TM</sup> is essential for fluorescence microscopy, confocal microscopic analysis or extracting during routine dehydration procedures? Most fluorescencelabeled samples are directly mounted in glycerol-based mounting media for fluorescence and confocal imaging. Under such conditions, a biological structure can be viewed at about 100µm beneath the surface by using laser source with wavelength at visible spectrum range. To increase depth of view, one can use two-photon confocal system with infrared laser as light source and viewing structures twice deeper than the conventional confocal microscope.  $FocusClear^{TM}$  is a water-soluble clearing agent for increases the transparency of biological tissues. As a result, image quality of fluorescence or non-fluorescence labeled specimens is greatly improved. FocusClear<sup>TM</sup> facilitates light penetration and allows visualization of internal objects up to at least 500µm below tissue surface. In contrast, traditional clearing agents such as glycerol-based mounting media allow visualization of only up to 150µm inside the specimen. Because of high tissue transparency, FocusClear<sup>TM</sup> also increases the efficacy of laser excitation and optical signal detection of either color or fluorescence.  $FocusClear^{TM}$  is suitable for microscopic observation of immunofluorescence

-labeled single cells and tissues, in situ hybridization, tissue or whole-mount immunohistochemistry, and fluorescence protein samples.

**Properties:** FocusClear<sup>TM</sup> solution is a water-soluble clearing agent. It is not gelling in the bottle and no dehydration of the objects is necessary. Samples can be directly transferred from water, buffer solutions, alcohol, DMSO, DMF, and glycerin into FocusClear<sup>TM</sup> solution. FocusClear<sup>TM</sup> can be used for samples labeled with fluorescence and non-fluorescence dyes including lipophilic dyes, such as DiI, DiD and NBD-ceramide. FocusClear<sup>TM</sup> is non-toxic, ready to use, always liquid, no need to be aliquoted, mixed, centrifuged or kept frozen. It allows easy and universal production of preparations.

 $MountClear^{TM}$  is a mountant which specially designed for

mounting specimens cleared by the  $FocusClear^{TM}$ . MountClear<sup>TM</sup> does not interfere the clearing effect of  $FocusClear^{TM}$ . In addition, it has anti-quenching,

non-fluorescence and quick clotting characteristics. Using mounting media other than  $MountClear^{TM}$  may result in cloudiness of the sample.

*Immersion Solution-M* is an immersion solution with a refraction index matching to those of *MountClear<sup>TM</sup>*. They are designed to avoid deformation of the observed images during high-resolution microscopic observation that using glycerin or water immersion objective lens.

**Effects:** Tissues in the *FocusClear<sup>TM</sup>* become transparent. The resolution and depth of focus greatly increased with sharp outline and high contrast. In order to obtain best results, it is recommended that the specimen cleared in *FocusClear<sup>TM</sup>* should be mounted in *MountClear<sup>TM</sup>* and observed with oil or water immersion lens with high numerical aperture and covered with *Immersion Solution-M. FocusClear<sup>TM</sup>*, however, is designed to clear specimens fixed by cross-linking agents such as paraformaldehyde and glutaraldehyde, but it is ineffective for heat-denatured or alcohol fixed specimens.

**Applicable tissues:** Mouse brain, insect brain, human tumors, some plants tissues and biomaterials.

#### 2. STORAGE AND HANDLING

Store at 4°C. Do not freeze. If slight turbidity occurs upon prolonged storage, clarify the solution by incubating in hot-water bath and centrifugation should be followed before use.

#### **3. WARNING AND PRECAUTIONS**

These products are intended for research purposes only. They may contain materials that are toxic to humans and animals and should not be administered either externally or internally.

#### **4. APPLICATION PROTOCOL**

1. Paraformaldehyde and/or glutaraldehyde fixed samples labeled with immunofluorescence, fluorescence probes, immunohistochemicals, or conventional dyes should be thoroughly washed to remove non-specific bindings.

2. Tissue blocks, brain slices, cryosections or single cells ready for microscopic observation can be directly transferred into appropriate amount of *FocusClear<sup>TM</sup>* solution for clearing.

Note: For an intact fly brain, 100µl FocusClear<sup>TM</sup> solution is suggested. For a slice of mouse brain (200 µm thick), 1 ml *FocusClear<sup>TM</sup>* solution is suggested.

3. For an effective clearing, the incubation time (10 min to 4 h) should be adjusted according to the size of the tissue ( $10^6 \ \mu m^3 \sim 1 \ mm^3$ ). To prevent evaporation during clearing, the incubation chamber should be completely sealed with parafilm membrane.

Note: Small samples such as fly brains may become completely transparent and difficult to be retrieved under a dissecting microscope. By simply applying a drop of saline solution, your precious samples will become visible again. You can clear the tissue again in a smaller drop of *FocusClear*<sup>TM</sup> for easy retrieval.

- 4. The cleared specimens are then mounted in a fresh drop of *FocusClear*<sup>TM</sup> solution.
- 5. For the best quality, the cleared specimens should be mounted in the *MountClear<sup>TM</sup>* solution. Prior to every use,

the *MountClear*<sup>TM</sup> solution should be completely dissolved again by incubation in the hot-water bath (55°C) for about 30 min. After brief cooling at room temperature, the *MountClear*<sup>TM</sup> solution is ready for use.

- 6. Seal the sample completely with fingernail polisher.
- 7. When using an oil/water immersion lens to observe the sample, *Immersion Solution-M* matching the reflective index of the mounting solution should be used for better resolution.

#### **5. TECHNICAL ASSITANCE**

E-mail: sales@celexplorer.com Mail: Technical Assistance Department, CelExplorer Labs. 7F-4, No 246, Sec. 2, Guangfu Rd., East Dist., HSINCHU 30099, TAIWAN(R.O.C.)

#### 6. REFERENCES

- 1. J. Comp. Neurol. (1999) 413, 593-602.
- 2. J. Comp. Neurol. (2001) 440, 1-11.
- 3. Proc. Natl. Acad. Sci. (2002) 99, 37-42.
- 4. Gastroenterology. (2009) 137(2), 453-65.

Item	Description	Cat#
FocusClear <sup>TM</sup> Kit	Includes 5 ml each of FC-101,MC-301, IS-502	F101-KIT
	Includes 50 ml each ofFC-102, MC-302, IS-503	F102-KIT
FocusClear <sup>TM</sup>	5 ml	FC-101
	50 ml	FC-102
<i>MountClear</i> <sup>TM</sup>	5 ml	MC-301
	50 ml	MC-302
Immersion Solution-M	5 ml	IS-502
	50 ml	IS-503

### PRODUCT LIST

CelExplorer's products are high-quality biochemical reagents for laboratory use only. These reagents are not for drug and medical purposes, household or other uses. Most CelExplorer's products and product applications are covered by U.S. and international patents.