PRODUCT INFORMATION



FGTA

Item No. 33963

CAS Registry No.: Formal Name:	67-42-5 3,12- <i>bis</i> (carboxymethyl)-6,9-dioxa-3,12- diazatetradecanedioic acid
Synonyms:	Egtazic Acid, Ethylene Glycol <i>bis</i> (β- aminoethyl ether)-N,N,N',N'-tetraacetic Acid, Ethylene Glycol Tetraacetic Acid, GEDTA, NSC 615010
MF:	$C_{14}H_{24}N_2O_{10}$ HO 0
FW:	380.4
Supplied as:	A solid
Storage:	-20°C
Stability:	≥2 years
1 6 1	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

EGTA is supplied as a solid. Aqueous solutions of EGTA can be prepared by directly dissolving the solid in aqueous buffers. EGTA is slightly soluble in PBS (pH 7.2). We do not recommend storing the aqueous solution for more than one day.

Description

EGTA is an aminopolycarboxylic acid chelating agent.¹ It is selective for calcium over magnesium with K_{app} values for calcium ranging from 6.93 to 10.7 x 10⁵ M⁻¹ in various buffers at pH 6.8 and 20°C. EGTA (5 mM) inhibits apoptosis induced by the sarcoplasmic/endoplasmic Ca²⁺-ATPase (SERCA) inhibitor thapsigargin (Item No. 10522) in isolated rat thymocytes.² It has been used as an irrigation agent to remove the smear layer from the pulp canal wall on extracted human teeth.³ Intranasal administration of EGTA enhances adenovirus-mediated gene transfer of LacZ to the trachea and main bronchi in mice by 8-fold.⁴

References

- 1. Harafuji, H. and Ogawa, Y. Re-examination of the apparent binding constant of ethylene glycol bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid with calcium around neutral pH. J. Biochem. 87(5), 1305-1312 (1980).
- 2. Jiang, S., Chow, S.C., Nicotera, P., et al. Intracellular Ca²⁺ signals activate apoptosis in thymocytes: Studies using the Ca²⁺-ATPase inhibitor thapsigargin. Exp. Cell. Res. 212(1), 84-92 (1994).
- 3. Viswanath, D., Hedge, A.M., and Munshi, A.K. The removal of the smear layer using EGTA: A scanning electron microscopic study. J. Clin. Pediatr. Dent. 28(1), 69-74 (2003).
- 4. Gregory, L.G., Harbottle, R.P., Lawrence, L., et al. Enhancement of adenovirus-mediated gene transfer to the airways by DEAE dextran and sodium caprate in vivo. Mol. Ther. 7(1), 19-26 (2003).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/09/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM