




Synthetic Auxin and Selective Binding to Modified Receptor



Synthetic auxin cvxIAA selectively binds to the modified auxin receptor, allowing for targeted binding at selected tissues and organs expressing the modified receptor.

Due to its multiple physiological functions, it has been a challenge to analyze several physiological phenomena stemming from auxin signaling at specific tissues and organs. Recently, Torii et al. at the Institute of Transformative Bio-Molecules (ITbM), at Nagoya University developed cvxIAA [M3141], a synthetic auxin using a molecular design technology known as the bump-and-hole strategy*. cvxIAA specifically binds to modified TIR1 receptor proteins (concave receptor, ccvTIR1). cvxIAA scarcely binds to wild type of TIR1 receptor protein and endogenous auxin scarcely binds to concave receptor. This cross-over independence and selective binding observed with Auxin-TIR1 versus cvxIAA-TIR1(modified) can allow for transformative research into otherwise difficult to study organ and tissue systems.

[Pairing of convex IAA (cvxIAA) and concave TIR1 (ccvTIR1) engineered using bump-and-hole strategy]



*The bump-and-hole strategy is a method used to engineer an orthogonal pair of receptor proteins and ligands by producing a small hole at a binding site in a receptor through mutation then subsequently introducing modified potential ligands to the binding site of the receptor. For the cvxIAA-ccvTIR1 system, modified TIR1 receptor proteins (concave TIR1, ccvTIR1) feature an amino acid substitution in the binding site with that lacks aromaticity. Alternatively, cvxIAA is designed to bind specifically to the modified TIR1 receptor protein by introducing an aromatic ring to structure of IAA.

Synthetic Auxin and Selective Binding to Modified Receptor

Application

The comparison of auxin (1-NAA) and convex IAA (cvxIAA) responses in seedlings of Arabidopsis thaliana wild type and transformants expressing concave receptors.

Chemical hijacking of auxin signaling with an engineered auxin-TIR1 pair N. Uchida, K. Takahashi, R. Iwasaki, R. Yamada, M. Yoshimura, T. A. Endo, S. Kimura, H. Zhang, M. Nomoto, Y. Tada, T. Kinoshita, K. Itami, S. Hagihara, K. U. Torii, Nat. Chem. Biol. 2018, 14, 299.

These information have been provided by Prof. Keiko Torii at Institute of TransformativeBio-Molecules (ITbM), Nagoya University.

Other auxins

IAA	5g / 25g [10022]	4-BPA	5g / 25g [B2746]
K-IAA	1g / 25g [I0023]	MCPA	25g / 500g [C0206]
IAA Methyl Ester	5g / 25g [M2605]	4-CPA	25g / 500g [C0250]
IAA Ethyl Ester	5g / 25g [E0878]	PCIB	25g [C0940]
IPA	5g / 25g [10032]	Dichlorprop	5g / 25g [D1942]
IBA	5g / 25g [10026]	Dicamba	25g [D4800]
IAN	1g / 25g [10024]	MCPB Ethyl Ester	100mg / 1g [E1149]
NAA	25g / 500g [N0005]	2,4-D	25g / 500g [D0396]
K-NAA	25g [N0006]	Na-2,4-D Monohydrate	25g / 500g [D1319]
Na-NAA	25g / 500g [N000 7]	2,4,5-T Potassium Salt	25g / 500g [T1509]
NOA	25g / 500g [N0045]		

1-Naphthaleneacetamide

25g / 500g [N0624]





Ordering and Customer Service

TCI AMERICA

:800-423-8616 / 503-283-1681 :888-520-1075 / 503-283-1987 E-mail: Sales-US@TCIchemicals.com

TCI EUROPE N.V.

: +32 (0)3 735 07 00 Fax : +32 (0)3 735 07 01 E-mail: Sales-EU@TCIchemicals.com

TCI Deutschland GmbH

: +49 (0)6196 64053-00 Fax : +49 (0)6196 64053-01 E-mail: Sales-DE@TCIchemicals.com

Tokyo Chemical Industry UK Ltd.

: +44 (0)1865 784560 Fax : +44 (0)1865 784561 E-mail: Sales-UK@TCIchemicals.com TCI Chemicals (India) Pvt. Ltd.

: 1800 425 7889 / 044-2262 0909 Tel Fax : 044-2262 8902

E-mail: Sales-IN@TCIchemicals.com

梯希爱(上海)化成工业发展有限公司 :800-988-0390 / 021-67121386

: 021-6712-1385 E-mail: Sales-CN@TCIchemicals.com

TOKYO CHEMICAL INDUSTRY CO., LTD.

Tel : +81 (0)3-5640-8878 : +81 (0)3-5640-8902 Fax

E-mail: global business@TCI chemicals.com

Availability, price or specification of the listed products are subject to change without prior notice. Reproduction forbidden without the prior written consent of Tokyo Chemical Industry Co., Ltd.