

PRODUCT INFORMATION



Ferulic Acid methyl ester

Item No. 17881

CAS Registry No.: 2309-07-1

Formal Name: 3-(4-hydroxy-3-methoxyphenyl)-2-propenoic acid,
methyl ester

Synonyms: Methyl Ferulate,
Methyl 4'-hydroxy-3'-methoxycinnamate

MF: C₁₁H₁₂O₄

FW: 208.2


Purity: ≥98%

UV/Vis.: λ_{max}: 219, 237, 326 nm

Supplied as: A crystalline solid

Storage: -20°C

Stability: As supplied, 2 years from the QC date provided on the Certificate of Analysis, when stored properly



Laboratory Procedures

Ferulic acid methyl ester is supplied as a crystalline solid. A stock solution may be made by dissolving the ferulic acid methyl ester in the solvent of choice. Ferulic acid methyl ester is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of ferulic acid methyl ester in ethanol and DMF is approximately 20 mg/ml and approximately 10 mg/ml in DMSO.

Ferulic acid methyl ester is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, ferulic acid methyl ester should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Ferulic acid methyl ester has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Ferulic acid is a hydroxycinnamic acid that is abundant in plants and originally derived from giant fennel (*F. communis*). This naturally-occurring phenolic has antioxidant activities that provide protection against inflammation and cancer.¹⁻⁴ Ferulic acid methyl ester is a lipophilic derivative of ferulic acid, demonstrating increased ability to cross cell membranes.⁵ Ferulic acid methyl ester has less antioxidant capacity than ferulic acid in neuronal PC12 cells (IC₅₀ = 74.7 μM for ferulic acid ethyl ester vs. 44.6 μM for ferulic acid, 2,2-diphenyl-1-picrylhydrazyl radical scavenging).⁶ Ferulic acid methyl ester, at 10-25 μg/ml, inhibits the release of pro-inflammatory cytokines, blocks the expression of COX-2, and reduces nitric oxide generation from LPS-stimulated macrophages.⁷

References

1. Huang, M.T., Smart, R.C., Wong, C., et al. *Cancer Res.* **48**, 5941-5946 (1988).
2. Conney, A.H., Lysz, T., Ferraro, T., et al. *Adv. Enzyme Regul.* **31**, 385-396 (1991).
3. Iwahashi, H., Nishizaki, K., and Takagi, I. *Biochem. J.* **361**, 57-66 (2002).
4. Sudheer, A.R., Muthukumaran, S., Devipriya, N.A., et al. *Toxicology* **243**, 317-329 (2008).
5. Sultana, R. *Biochim. Biophys. Acta* **1822(5)**, 748-752 (2012).
6. Garrido, J., Gaspar, A., Garrido, E.M., et al. *Biochimie* **94(4)**, 961-967 (2012).
7. Phuong, N.T.M., Cuong, T.T., and Quang, D.N. *Asian Pac. J. Trop. Med.* **7(Suppl 1)**, S327-S331 (2014).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/12/2016

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM