

## **PAX5** Polyclonal Antibody

Catalog No.	A3068
Applications	WB
Cross-reactivity	Human

Category **Observed MW** 

Calculated MW

**Polyclonal Antibodies** 42kDa 31kDa/32kDa/34kDa/35kDa/37kDa/38kDa/42kDa

1:3000


## **Immunogen Information**

Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 150-300 of human PAX5 (NP_057953.1).
Gene ID	5079
Swiss prot	Q02548
Synonyms	PAX5; ALL3; BSAP; paired box 5
Product information	

Source	Rabbit
Isotype	lgG
<b>Purification method</b>	Affinity purification
Storage	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

## Background

This gene encodes a member of the paired box (PAX) family of transcription factors. The central feature of this gene family is a novel, highly conserved DNA-binding motif, known as the paired box. Paired box transcription factors are important regulators in early development, and alterations in the expression of their genes are thought to contribute to neoplastic transformation. This gene encodes the B-cell lineage specific activator protein that is expressed at early, but not late stages of B-cell differentiation. Its expression has also been detected in developing CNS and testis and so the encoded protein may also play a role in neural development and spermatogenesis. This gene is located at 9p13, which is involved in t(9;14)(p13;q32) translocations recurring in small lymphocytic lymphomas of the plasmacytoid subtype, and in derived large-cell lymphomas. This translocation brings the potent E-mu enhancer of the IgH gene into close proximity of the PAX5 promoter, suggesting that the deregulation of transcription of this gene contributes to the pathogenesis of these lymphomas. Alternative splicing results in multiple transcript variants encoding different isoforms.



**Recommended Dilutions** 

